

Running head: Probiotics in COVID-19

Title: Probiotics and COVID-19: is there any link?

Amal Akour^{1,2}, BPharm., PhD.

¹Department of Biopharmaceutics and Clinical Pharmacy, The School of Pharmacy, The University of Jordan, Jordan

²Department of Pharmacy, Faculty of Pharmacy, Al-Zaytoonah University of Jordan, Jordan

Address: Queen Rania Street, Amman 11942 Jordan

Email: a.akour@ju.edu.jo, amalakour@gmail.com

Significance and Impact of Study

The role of probiotics in alleviation of the novel COVID-19 has not been established. This review provides an insight about the anti-inflammatory and antiviral effects of probiotics *in vitro*, animal models and human. The latter, can provide an indirect evidence and/or hypothesis-driven approach to investigate the use probiotics as adjunctive therapy in the prophylaxis and/or alleviation of COVID-19 symptoms.

ABSTRACT

Understanding mechanisms of the novel SARS-CoV2 infection and progression, can provide potential novel targets for prevention and/or treatment. This could be achieved via the inhibition of viral entry and/or replication, or by suppression the immunologic response that is provoked by the infection (known as the cytokine storm). Probiotics are defined as “live microorganisms that, when administered in adequate amounts, confer a health benefit on the host”. There is scarcity of evidence about the relationship between COVID-19 and gut microbiota. So, whether or not these supplements can prevent or ameliorate COVID-19 associated symptoms is not fully understood. The aim of this paper is to provide an indirect evidence about the utility of probiotics in combating COVID-19 or its associated symptoms, through the review of its antiviral and anti-inflammatory properties *in vitro*, animal models, and human trials.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/LAM.13334](#)

This article is protected by copyright. All rights reserved

KEYWORDS: COVID-19, probiotics, anti-inflammatory, antiviral, adjunctive therapy, microbiota

INTRODUCTION

The novel coronavirus SARS-CoV-2 pandemic has emerged in late December of 2019 (Eurosurveillance Editorial, 2020). To date, there is no established prevention or treatment protocol for this virus. Treatments are based on investigational repurposed drugs as well as symptoms alleviation. Small molecules that have shown to prevent viral entry or replication such as hydroxychloroquine (Yazdany and Kim, 2020) and remdesivir (Cao et al., 2020) are being studied in multiple clinical trials worldwide. Moreover, biologics such as tocilizumab and interferon beta are being tested for their ability to suppress inflammation and the so called “the cytokine storm” (Liu et al., 2020). Indeed, COVID-19 patients who have had higher concentrations of proinflammatory cytokines and chemokines, were more likely to be admitted to an Intensive care unit, namely, Granulocyte-colony stimulating factor (G-CSF), The human interferon-inducible protein 10 (IP-10/CXCL10), monocyte chemoattractant protein-1 (MCP-1), and tumor necrosis factor-alpha (TNF- α), in addition to elevated cytokines from T helper 2 cells such as Interleukin (IL)-4 and, IL-10 (Liu et al., 2020).

According to World Health Organization (WHO), Food Agriculture Organization (FAO), and the International Scientific Association for Probiotics and Prebiotics (ISAPP), Probiotics are defined as “live microorganisms that, when administered in adequate amounts, confer a health benefit on the host” (Hill et al., 2014). The most commonly used microorganisms are *Lactobacillus spp.*, *Bifidobacteria* and *Saccharomyces* (Zendeboodi et al., 2020; Guimarães et al., 2020). They are available commercially as add-on to several food and dairy products, and supplements (de Almada et al., 2015; Kiousi et al., 2019; Roobab et al., 2020). Thereafter, the definition of the term probiotic has been modified and extended. Although, the definition developed by WHO and FAO is generally conventional, new terminologies have emerged lately in the literature such as ‘paraprobiotics’ (dead/inactivated cells of probiotics) and ‘postbiotics’ (healthful metabolites of probiotics) (Zendeboodi et al., 2020; Barros et al., 2020). In addition, there are prebiotics which are “substrates that are selectively utilized by host microorganisms conferring a health benefit” (Neri-Numa et al., 2020). Most of prebiotics are carbohydrate-based, but that phenolic compounds and conjugated fatty

acids can comply with the criteria of prebiotics as well. Prebiotics have also shown many health effects such as pathogen inhibitory effects and immune modulation, etc. (Neri-Numa et al., 2020). However, for the sake of this review will concentrate of probiotics.

It is well established that probiotics, mainly the strains of lactic acid bacteria, can modulate the human gut microbiota via the suppression of opportunistic bacteria growth (Khaneghaha et al., 2020). Therefore, the administration as well as the stimulation of the growth and activity of probiotic strains in gut can be considered as a potential approach to control foodborne enteric pathogens. Moreover, there many health benefits of probiotics beyond the gut, probiotics have shown to enhance the immunity, reduce the severity of certain allergic conditions, in addition to conferring some anti-carcinogenic properties (Khaneghaha et al., 2020). There are scarce data available about the effect of COVID-19 on intestinal microbiota. A recent small study from China showed that patients with COVID-19 showed microbial dysbiosis with decreased *Lactobacillus* and *Bifidobacterium* counts (Xu et al., 2020). The same study proposed using prebiotics or probiotics as adjunctive therapy to regulate the balance of intestinal microbiota and reduce the risk of secondary infection in those patients. On the other side, animal studies (though not peer-reviewed yet) indicated that *Lactobacillus acidophilus* and *Bacillus clausii* did not reduce coronavirus receptor expression in the murine small intestine compared with control and post-Salmonella infection models (Feng et al., 2020). It was reported that 58–71% of patients with COVID-19 in China were given antibiotics, and antibiotic-related diarrhea occurred in 2–36% of patients (Mak et al., 2020). In those cases, supplementing the gut normal flora by using probiotics has been proposed make COVID-19 patients less prone to secondary infections.

Human gut microbiota is a very diverse system and we should examine the efficacy other types of beneficial bacteria, not only *Lactobacilli* and *Bifidobacteria*, in fighting COVID- 19 or alleviation of the associated inflammation. To date, the rationale for using probiotics in COVID-19 patients can be merely extrapolated from an indirect hypothetical evidence (Mak et al., 2020;Baud et al., 2020). Therefore, we cannot draw any conclusions unless we have well-designed preclinical and clinical studies.

Of interest, studies have shown that probiotics can have potential anti-inflammatory and anti-viral effects. Indeed, they act by suppressing cytokines production whether locally, i.e. on the intestinal

mucosa level, or on extra-intestinal body organs (Kiouisi et al., 2019). Several clinical trials have shown that the administration of probiotics to improve outcomes in immune system-related conditions, or viral infections, such as atopic dermatitis, rheumatoid arthritis, and some allergic conditions, or respiratory tract infections (Kiouisi et al., 2019).

Based on the above, it is plausible to assume that these anti-inflammatory and antiviral effects could potentially contribute, at least partially or in combination with other medications, in prevention and/or alleviation of COVID-19-related symptoms.

RESULTS AND DISCUSSION

Antiviral activity

It was shown that probiotics in adults or children was safe (Tapiovaara et al., 2016) and clinically effective (Wang et al., 2016) in reducing the severity and the duration of upper respiratory tract infections. These were confirmed via a review (Kanauchi et al., 2018) where the ability of different strains lactic acid-producing bacteria to ameliorate, or prevent different viral infections was illustrated, such as decreasing titers of Ebola and cytomegalovirus, decreasing severity and duration of upper respiratory tract infection or gastroenteritis. The main focus of that review, however, was the anti-influenza activity of *L. lactis* JCM 5805, which has the potency to directly stimulate plasmacytoid dendritic cells (pDCs) via TLR9, which will in turn stimulate interferon production and control viral replication and spread. The activity of probiotic strain *B. subtilis* (Starosila et al., 2017) against the influenza virus *in vitro* and in animals was investigated. A new peptide, P18, produced by the probiotic strain was isolated, characterized, and synthesized thereafter. Interestingly, while being nontoxic, P18 exhibited a complete inhibition of the influenza virus *in vitro*. Moreover, the antiviral effect of P18 in mice was comparable to that of oseltamivir phosphate (Tamiflu®). Recently, the effect *Lactobacillus gasseri* in prophylaxis against respiratory syncytial virus (RSV) infection was evaluated in mice (Eguchi et al., 2019). The RSV titer in the lung was significantly decreased, and the expression of pro-inflammatory cytokines in the lung were significantly diminished while interferons and interferon stimulated genes were upregulated after the treatment. In clinical settings, a randomized, double-blind, placebo-controlled trial have shown that the administration of prebiotics and probiotics early in life of preterm infants resulted in a significant

lower the incidence of viral respiratory tract infections, especially those caused by rhinovirus, even though viral RNA load was the same in the probiotic and the placebo groups (Luoto et al., 2014). Based on the above data, the alteration of gut microbiota with good choice of prebiotics and probiotics might offer a novel and cost-effective methodologies to reduce the risk of viral infections.

Anti-inflammatory activity

In vitro/ex vivo studies

A recent paper (Yu et al., 2019), evaluated the anti-inflammatory potential of *Weissella cibaria* (JW15) against lipopolysaccharide (LPS) stimulation in RAW 264.7 cells. Treatment with heat-killed JW15 decreased nitric oxide and prostaglandin E2 production via downregulation of the inducible nitric oxide synthase and cyclooxygenase-2. In addition, treatment with heat-killed JW15 suppressed the expression of pro-inflammatory cytokines, IL-1 β , IL-6, and TNF- α . The anti-inflammatory properties of treating with heat-killed JW15 were associated with mitogen-activated protein kinase signaling pathway mediated suppression of nuclear factor- κ B (NF- κ B). These results indicated that JW15 possesses anti-inflammatory potential and provide a molecular basis regarding the development of functional probiotic products.

Moreover, the effect of potential probiotic strains on early gingivitis was evaluated in an ex vivo model (Schmitter et al., 2018). Human monocytes was stimulated by bacterial LPS. Strains *Lactobacillus paracasei* and *Lactobacillus plantarum* induced statistically significant and dose-dependent reductions in the release of multiple inflammatory mediators including IL-6, IL-8 and prostaglandin E2 vs. placebo.

The effects of two strains of *L. rhamnosus*, *L. rhamnosus GG* and *GR-1*, in modulating production of TNF- α in human monocytic cell line THP-1 and mouse macrophages was investigated. (Kim et al., 2006). In C57BL/6 mouse bone marrow-derived immortalized macrophages (BMDIM), and THP-1 cells, both strains suppressed the production of *E.coli*- and LPS-induced TNF α production. This in vitro study showed that this suppression was likely mediated through downregulation of TNF- α mRNA expression and a paracrine effect of G-CSF, as G-CSFR $^{-/-}$ macrophages produced similar amounts of TNF- α as wild-type cells. G-CSF induced the activation of Signal Transducer And

Activator Of Transcription 3 (STAT3) protein, which in turn prevented JunN-terminal kinases (JNK) activation and consequently resulted in suppression of TNF- α production in LPS- or E. coli-activated macrophages. A comprehensive review (Plaza-Diaz et al., 2017) summarized five *in vitro* studies which assessed the anti-inflammatory role of antibiotics in intestinal diseases. Pretreatment of Caco2 cells or dendritic cells with several probiotics decreased production of several inflammatory chemokines and cytokines, such as IL-22, IL-8 and TNF- α , and downregulation of mitogen-activated protein kinase (MAPK) and NF- κ B pathway or even ameliorated enterocyte apoptosis in intestinal epithelial cells.

Animal models

Two strains of *Lactobacillus* (*mucosae* and *fermentum*) were assessed for anti-inflammatory properties (Ayyanna et al., 2018) in acute and chronic Freund's adjuvant-induced inflammation model. The two probiotic strains were administered orally along with feed to the Wistar albino male rats as whole cell as well as microencapsulated form. Both forms of the probiotics lead to a significant decrease in paw edema. Percentage of inhibition in paw thickness of microencapsulated probiotic bacteria, un-encapsulated strains were revealed $85 \pm 13\%$ and $77 \pm 25\%$, respectively. In addition, there was an upregulation of anti-inflammatory cytokine genes and levels (IL-10) and down-regulation pro-inflammatory cytokine genes and levels (IL-6 and TNF- α) in probiotic-treated rat. The paw tissues of the probiotic-treated rats have also exhibited the low level of lipid peroxides formation and higher anti-oxidant activities when compared to the control. The anti-inflammatory role of probiotics in various animal models in different forms of colitis was reviewed (Plaza-Diaz et al., 2017), and it was found that the administration of probiotics ameliorated these conditions as evidenced through histological improvement, reduction of disease severity index or decline in the levels of inflammatory markers such as myeloperoxidase (MPO) activity, nitric oxide synthase (iNOS), cyclooxygenase-2 (COX-2), NF- κ B, TNF α , IL-6, and phosphorylated Akt, while increasing IL-10 expression in colonic tissue (Plaza-Diaz et al., 2017). In animal models of multiple sclerosis, (Morshedi et al., 2019), it was shown that giving probiotics reduced T-cell autoreactive cell response, production of pro-inflammatory cytokines and increased levels on anti-inflammatory markers such as IL-10 and nitric oxide.

Human studies

In two clinical studies, the administration of probiotics resulted in reduction of IL-6, C-reactive protein (CRP), while there was increased levels of IL-10 in the sera of the multiple sclerosis patients (Morshedi et al., 2019). One clinical study (Plaza-Diaz et al., 2017) showed that probiotics administration reduced the systemic pro-inflammatory biomarkers in both gastrointestinal and non-gastrointestinal conditions in colitis patients after 6-8 weeks of treatment.

The functional properties of 8 *Lactobacillus* strains from infant, which showed probiotic potential as well as ability to resist acidic pH and bile salts (Oh et al., 2018) These isolate included various *Lactobacillus* species: *L. reuteri* 3M02 and 3M03, *L. gasseri* 4M13, 4R22, 5R01, 5R02, and 5R13, and *L. rhamnosus* 4B15, which were evaluated for anti-oxidation, inhibition of α -glucosidase activity, cholesterol-lowering, and anti-inflammatory activity. Particularly, two strains of *L. rhamnosus*, 4B15 (4B15) and *L. gasseri* 4M13 (4M13) showed considerably higher anti-oxidation, inhibition of α -glucosidase activity, and cholesterol-lowering, and greater inhibition of nitric oxide production than other strains. Moreover, the aforementioned strains substantially inhibited the release of inflammatory mediators such as TNF α , IL-6, IL-1 β , but increased IL-10 in LPS-stimulated RAW 264.7 macrophages.

Regarding the effect of probiotics in respiratory illnesses, two randomized controlled trials showed that critically ill patients on mechanical ventilation who were administered probiotics (*Lactobacillus rhamnosus* GG, live *Bacillus subtilis*, and *Enterococcus faecalis*) had significantly less ventilator-associated pneumonia compared with placebo (Morrow et al., 2010; Zeng et al., 2016). Accordingly, it can be speculated that COVID-19-related pneumonia could be alleviated in the same manner.

No published studies so far evaluated the use of probiotics as add-on therapy for the management of COVID-19, so their role is not yet established. . However, according a recent science blog by IASPP, many researchers all over the world are studying the relationship between microbiome and susceptibility to COVID-19 as well as assessing the ability of various probiotics strains to reduce the viral load via different mechanisms of action (International Scientific Association of Probiotics and Prebiotics board of directors, 2020). Currently a research team in Belgium, is exploring the potential of specific strains of *Lactobacilli* in the nasopharynx and oropharynx to reduce viral activity via the

barrier-enhancing and anti-inflammatory effects, and reduce the risk of secondary bacterial infections in COVID-19 (International Scientific Association of Probiotics and Prebiotics board of directors, 2020). In the United Kingdom, there is an ongoing Phase II randomized, double-blind, placebo-controlled trial to evaluate the efficacy and safety of oral Live Biotherapeutic MRx-4DP0004 (2 capsules) in addition to standard supportive care for hospitalized COVID-19 patients for 14 days. In the USA, a multicenter, randomized, double blind, phase 2 trial using a commercial product of *Lactobacillus rhamnosus* GG (20 billion CFU) vs. placebo are conducted to see its ability to decrease infections and improve outcomes. The same team is in the process of developing protocols to study prevention and treatment of COVID-19 in a range of other at-risk populations including hospitalized patients and healthcare workers.

In lieu of their antiviral and anti-inflammatory activity, although very early to judge, it is tempting to assume that probiotics can potentially be among the rational adjunctive options for the treatment and prophylaxis of viral infections including COVID-19 infection. In a preprint paper by Feng *et al* (Feng et al., 2020), RNA sequencing results showed that the coronavirus receptors, including angiotensin-converting enzyme 2 (ACE2) for SARS-CoV and SARS-CoV2, were highly expressed in human enterocytes. Interestingly, these potential target cells have constant expression in the small intestine is constant while being continuously changed in lung tissues. Therefore, enterocytes may act as a conserved cell reservoir for coronaviruses, a fact that should draw the attention of health researchers to this site of SARS-CoV2 infection.

In conclusion, until there is more evidence about the novel coronavirus pathogenesis and its effect on gut microbiome, the use of conventional probiotics for COVID-19 cannot be liberally recommended. Although gut dysbiosis has been suggested in the pathogenesis of certain respiratory conditions (Li et al., 2021), more targeted and novel approaches are warranted. Although, the modulation of gut microbiota is expected to be one of the promising therapeutic approaches to alleviate COVID-19 and/or its associated inflammatory complications, still, the utility of probiotics as add-on therapy may need to be further studied in well-designed randomized controlled clinical trials to establish their efficacy and safety.

MATERIALS AND METHODS

Search was done using the following database including Google Scholar, PubMed using the following MeSH terms: COVID-19, probiotics, novel coronavirus, respiratory tract infection, anti-inflammatory, antiviral.

ACKNOWLEDGEMENTS. No funding was received no produce this work.

CONFLICT OF INTEREST. The author has no conflict of interest to declare.

References

- Ayyanna, R., Ankaiah, D. and Arul, V. (2018). Anti-inflammatory and Antioxidant Properties of Probiotic Bacterium *Lactobacillus mucosae* AN1 and *Lactobacillus fermentum* SNR1 in Wistar Albino Rats. *Front Microbiol* **9**, 3063.
- Barros, C. P., Guimarães, J. T., Esmerino, E. A., Duarte, M. C. K., Silva, M. C., Silva, R., Ferreira, B. M., Sant'ana, A. S., Freitas, M. Q. and Cruz, A. G. (2020). Paraprobiotics and postbiotics: concepts and potential applications in dairy products. *Current Opinion in Food Science* **32**, 1-8.
- Baud, D., Agri, V. D., Gibson, G. R., Reid, G. and Giannoni, E. (2020). Using Probiotics to Flatten the Curve of Coronavirus Disease COVID-2019 Pandemic. *Front. in Public Health* **8**, 186.
- Cao, Y. C., Deng, Q. X. and Dai, S. X. (2020). Remdesivir for severe acute respiratory syndrome coronavirus 2 causing COVID-19: An evaluation of the evidence. *Travel Med Infect Dis*, 101647.
- De Almada, C. N., Nunes De Almada, C., Martinez, R. C. and Sant'ana Ade, S. (2015). Characterization of the intestinal microbiota and its interaction with probiotics and health impacts. *Appl Microbiol Biotechnol* **99**, 4175-99.
- Eguchi, K., Fujitani, N., Nakagawa, H. and Miyazaki, T. (2019). Prevention of respiratory syncytial virus infection with probiotic lactic acid bacterium *Lactobacillus gasseri* SBT2055. *Sci Rep* **9**, 4812.
- Eurosurveillance Editorial, T. (2020). Latest updates on COVID-19 from the European Centre for Disease Prevention and Control. *Euro Surveill* **25**.
- Feng, Z., Wang, Y. and Qi, W. (2020). The small intestine, an underestimated site of SARS-CoV-2 infection: from Red Queen effect to probiotics. *Preprints*.
- Guimarães, J. T., Balthazar, C. F., Silva, R., Rocha, R. S., Graça, J. S., Esmerino, E. A., Silva, M. C., Sant'ana, A. S., Duarte, M. C. K., Freitas, M. Q. and Cruz, A. G. (2020). Impact of probiotics and prebiotics on food texture. *Current Opinion in Food Science* **33**, 38-40.

Hill, C., Guarner, F., Reid, G., Gibson, G. R., Merenstein, D. J., Pot, B., Morelli, L., Canani, R. B., Flint, H. J., Salminen, S., Calder, P. C. and Sanders, M. E. (2014). Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol* **11**, 506-14.

International Scientific Association of Probiotics and Prebiotics Board of Directors. 2020. *How some probiotic scientists are working to address COVID-19* [Online]. ISAPP. Available: <https://isappscience.org/how-some-probiotic-and-prebiotic-scientists-are-working-to-address-covid-19/> [Accessed 19/5/2020].

Kanauchi, O., Andoh, A., Abubakar, S. and Yamamoto, N. (2018). Probiotics and Paraprobiotics in Viral Infection: Clinical Application and Effects on the Innate and Acquired Immune Systems. *Curr Pharm Des* **24**, 710-717.

Khaneghaha, A. M., Abharib, K., Eş, I., Soaresa, M. B., Oliveiraa, R. B. A., Hosseinib, H., Rezaeie, M., Balthazara, C. F., Silva, R., Cruzd, A. G., Ranadheerah, S. and Sant'anaa, A. S. (2020). Interactions between probiotics and pathogenic microorganisms in hosts and foods: A review. *Trends in Food Science & Technology* **95**, 205-218.

Kim, S. O., Sheikh, H. I., Ha, S. D., Martins, A. and Reid, G. (2006). G-CSF-mediated inhibition of JNK is a key mechanism for *Lactobacillus rhamnosus*-induced suppression of TNF production in macrophages. *Cell Microbiol* **8**, 1958-71.

Kiousi, D. E., Karapetsas, A., Karolidou, K., Panayiotidis, M. I., Pappa, A. and Galanis, A. (2019). Probiotics in Extraintestinal Diseases: Current Trends and New Directions. *Nutrients* **11**.

Li, D., Zhao, M. Y. and Malcolm, T. H. (2021). What makes a foodborne virus: comparing coronaviruses with human noroviruses. *Current Opinion in Food Science* **42**, 1-7.

Liu, C., Zhou, Q., Li, Y., Garner, L. V., Watkins, S. P., Carter, L. J., Smoot, J., Gregg, A. C., Daniels, A. D., Jervey, S. and Albaiu, D. (2020). Research and Development on Therapeutic Agents and Vaccines for COVID-19 and Related Human Coronavirus Diseases. *ACS Cent Sci* **6**, 315-331.

Luoto, R., Ruuskanen, O., Waris, M., Kalliomaki, M., Salminen, S. and Isolauri, E. (2014). Prebiotic and probiotic supplementation prevents rhinovirus infections in preterm infants: a randomized, placebo-controlled trial. *J Allergy Clin Immunol* **133**, 405-13.

Mak, J. W. Y., Chan, F. K. L. and Ng, S. C. (2020). Probiotics and COVID-19: one size does not fit all. *The Lancet Gastroenterology and Hepatology* **10.1016/S2468-1253(20)30122-9**.

- Morrow, L. E., Kollef, M. H. and Casale, T. B. (2010). Probiotic prophylaxis of ventilator-associated pneumonia: a blinded, randomized, controlled trial. *Am J Respir Crit Care Med* **182**, 1058-64.
- Morshedi, M., Hashemi, R., Moazzen, S., Sahebkar, A. and Hosseiniard, E. S. (2019). Immunomodulatory and anti-inflammatory effects of probiotics in multiple sclerosis: a systematic review. *J Neuroinflammation* **16**, 231.
- Neri-Numa, A. I., Arruda, H. S., Geraldi, M. V., Maróstica Júnior, M. R. and Pastore, G. M. (2020). Natural prebiotic carbohydrates, carotenoids and flavonoids as ingredients in food systems. *Current Opinion in Food Science* **33**, 98-107.
- Oh, N. S., Joung, J. Y., Lee, J. Y. and Kim, Y. (2018). Probiotic and anti-inflammatory potential of *Lactobacillus rhamnosus* 4B15 and *Lactobacillus gasseri* 4M13 isolated from infant feces. *PLoS One* **13**, e0192021.
- Plaza-Diaz, J., Ruiz-Ojeda, F. J., Vilchez-Padial, L. M. and Gil, A. (2017). Evidence of the Anti-Inflammatory Effects of Probiotics and Synbiotics in Intestinal Chronic Diseases. *Nutrients* **9**.
- Roobab, U., Batool, Z., Manzoor, F. M., Asim, M., Moazzam, S., Khan, R. and Aadil, R. M. (2020). Sources, formulations, advanced delivery and health benefits of probiotics. *Current Opinion in Food Science* **32**, 17-28.
- Schmitter, T., Fiebich, B. L., Fischer, J. T., Gajfulin, M., Larsson, N., Rose, T. and Goetz, M. R. (2018). Ex vivo anti-inflammatory effects of probiotics for periodontal health. *J Oral Microbiol* **10**, 1502027.
- Starosila, D., Rybalko, S., Varbanetz, L., Ivanskaya, N. and Sorokulova, I. (2017). Anti-influenza Activity of a *Bacillus subtilis* Probiotic Strain. *Antimicrob Agents Chemother* **61**.
- Tapiovaara, L., Lehtoranta, L., Poussa, T., Makivuokko, H., Korpela, R. and Pitkaranta, A. (2016). Absence of adverse events in healthy individuals using probiotics--analysis of six randomised studies by one study group. *Benef Microbes* **7**, 161-9.
- Wang, Y., Li, X., Ge, T., Xiao, Y., Liao, Y., Cui, Y., Zhang, Y., Ho, W., Yu, G. and Zhang, T. (2016). Probiotics for prevention and treatment of respiratory tract infections in children: A systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)* **95**, e4509.
- Xu, K., Cai, H., Shen, Y., Ni, Q., Chen, Y., Hu, S., Li, J., Wang, H., Yu, L., Huang, H., Qiu, Y., Wei, G., Fang, Q., Zhou, J., Sheng, J., Liang, T. and Li, L. (2020). [Management of corona virus disease-19 (COVID-19): the Zhejiang experience]. *Zhejiang Da Xue Xue Bao Yi Xue Ban* **49**, 0.
- Yazdany, J. and Kim, A. H. J. (2020). Use of Hydroxychloroquine and Chloroquine During the COVID-19 Pandemic: What Every Clinician Should Know. *Ann Intern Med*.

Accepted Article

Yu, H. S., Lee, N. K., Choi, A. J., Choe, J. S., Bae, C. H. and Paik, H. D. (2019). Anti-Inflammatory Potential of Probiotic Strain *Weissella cibaria* JW15 Isolated from Kimchi through Regulation of NF-kappaB and MAPKs Pathways in LPS-Induced RAW 264.7 Cells. *J Microbiol Biotechnol* **29**, 1022-1032.

Zendeboodi, F., Khorshidian, N., Mortazavian, A. M. and Da Cruz, A. G. (2020). Probiotic: conceptualization from a new approach. *Current Opinion in Food Science* **32**, 103-123.

Zeng, J., Wang, C. T., Zhang, F. S., Qi, F., Wang, S. F., Ma, S., Wu, T. J., Tian, H., Tian, Z. T., Zhang, S. L., Qu, Y., Liu, L. Y., Li, Y. Z., Cui, S., Zhao, H. L., Du, Q. S., Ma, Z., Li, C. H., Li, Y., Si, M., Chu, Y. F., Meng, M., Ren, H. S., Zhang, J. C., Jiang, J. J., Ding, M. and Wang, Y. P. (2016). Effect of probiotics on the incidence of ventilator-associated pneumonia in critically ill patients: a randomized controlled multicenter trial. *Intensive Care Med* **42**, 1018-28.